

Available online at www.sciencedirect.com

SciVerse ScienceDirect

journal homepage: www.jfma-online.com

LETTER TO THE EDITOR

The application of urinary antigen test score in *Pneumococcal pneumonia* in children

Streptococcus pneumoniae is the most common pathogen of bacterial pneumonia in children.¹ The difficulty in acquiring good quality of sputum samples has been a major issue of pathogenic identification. Therefore, urine pneumococcal antigen test becomes a fast and feasible tool for diagnosis. However, high prevalence of pneumococcal colonization hinders its application in young children comparing to which in adult patients.² Shen et al³ proposed a urinary antigen test score to correlate with the severity of pneumococcal pneumonia in children. The results showed faster and greater enhancement of urinary antigen test was associated with severe symptoms. Although the report is inspiring, there are two issues to be addressed. First, the detection rate of urine pneumococcal antigen test increases in concentrated urine samples.⁴ In this study, the authors did not delineate the time when the urine antigen tests were performed and the concentration of each corresponding urine samples. It was difficult to clarify whether the faster and darker response of antigen test was due to the larger bacterial load in severe cases or was caused by more concentrated urine sample at the time point of collection. Second, the cases with underlying illness in this study apparently had diverse degree of severity. A case of cerebral palsy with impaired mobility may have greater chances predisposing pulmonary infection and experiencing more complicated course of illness compared with a child of diabetes mellitus with uneventful medical history. It would be better to spare the data of the cases with prior conditions from the study or to explain that the pulmonary function of these patients was in similar status. Nevertheless, the merit of

this study is proposing the relationship between bacterial load and the severity of pneumococcal pneumonia in children. Future efforts are required to pursue a more efficient and reliable diagnostic tool.

References

1. Michelow IC, Olsen K, Lozano J, Rollins NK, Duffy LB, Ziegler T, et al. Epidemiology and clinical characteristics of community-acquired pneumonia in hospitalized children. *Pediatrics* 2004; 113:701–7.
2. Dowell SF, Garman RL, Liu G, Levine OS, Yang YH. Evaluation of Binax NOW, an assay for the detection of pneumococcal antigen in urine samples, performed among pediatric patients. *Clin Infect Dis* 2001;32:824–5.
3. Shen CF, Wang SM, Liu CC. A new urinary antigen test score correlates with severity of pneumococcal pneumonia in children. *J Formos Med Assoc* 2011;110:613–8.
4. Dominguez J, Blanco S, Rodrigo C, Azuara M, Galí N, Mainou A, et al. Usefulness of urinary antigen detection by an immunochromatographic test for diagnosis of pneumococcal pneumonia in children. *J Clin Microbiol* 2003;41:2161–3.

Wen-Chan Huang
 Department of Pediatrics, National Taiwan University
 Hospital, College of Medicine, National Taiwan University,
 Taipei, Taiwan
 E-mail address: summerdebbie@gmail.com

9 September 2012